

Missouri Weekly Influenza Report 2006-2007 Season¹

Missouri is reporting "Sporadic Influenza Activity" to the CDC for Week 20².

To view influenza maps click here. Each map will give county data by placing the cursor over the county.

THIS IS THE FINAL REPORT FOR THE 2006-2007 SEASON.

Table 1. Reported Laboratory cases by sub-type for the Week ending May 19, 2007 (Week 20)

Serogroups	A (non- typed)	A (H1)	A (H1N1)	A (H3)	В	A or B Untyped (rapid test)	Total
Week 20	24	0	0	0	0	0	24

²⁴ cases by rapid non-culture diagnostic test for week 20.

Table 2. Influenza Season-to-Date and 5-season Median by Influenza Type Through Week Ending May 19, 2007 (Week 20)

Influenza Type	2006-07 Season	5-Season Median	Percent Change from 5-Season Median	
Influenza A	9,175	6,705	36.8%	
Influenza B	2,207	881	150.5%	
Influenza Unknown Or Untyped	3,463	3,052	13.5%	
Total	14,845	10,860	36.7%	

Season-to-Date: two isolates are identified as influenza A/Hawaii/15/2001-like (H1N1) viruses and two other isolates as influenza A/New Caledonia/20/99-like (H1N1) viruses.

Table 3. Influenza Season-to-Date and 5-season Median by Age Group Through Week Ending May 19, 2007 (Week 20)

Age Group	2006-07 Count	5-Season Median	Percent Change from 5-Season Median
00-<02	1,703	1,320	29.0%
02-04	2,503	991	152.6%
05-14	6,474	2,161	199.6%
15-24	1,323	1,134	16.7%
25-49	1,960	1,504	30.3%
50-64	528	599	(11.9%)
65+	354	1,275	(72.2%)
Total	14,845	10,860	36.7%

Table 4. Influenza Season-to-Date and 5-season Median by Region Through Week Ending May 19, 2007 (Week 20)

Region	2006-07 Count	5-Season Median	Percent Change from 5-Season Median
CE	1,875	1,598	17.3%
EA	3,643	4,146	(12.1%)
NW	3,856	2,653	45.3%
SE	1,786	680	162.6%
SW	3,685	1,587	132.2%
Total	14,845	10,860	36.7%

Table 5. Deaths involving Pneumonia and Influenza (P&I) Reported During the Week Ending May 12, 2007 (Week 19)*

Week 19	Season-to-Date	Week 19 Last Season	5 Year Weekly Median
60	2718	81	96

^{*} Beginning in Week 35 of 2003, the number of P&I deaths became based on a new system of retrieval that now includes <u>all</u> contributing causes of death from death certificates.

Graph 1. Influenza 2006-07 Season-To-Date as compared to the previous 4 influenza seasons Through the Week Ending May 19, 2007 (Week 20)




Table 6. U.S. Influenza Sentinel Physicians Surveillance Network Influenza-like Illness (ILI) for the Week Ending May 12, 2007 (Week 19)

Age	Age	Age	Age	Total ILI	Total	Percent
0-4	5-24	25-64	65+	Patients Seen	Patients Seen	ILI*
0	0	0	0	0	1267	0.00*

^{*}This is below the national baseline of 2.1%

Table 7. Respiratory Specimens Submitted to SPHL for Viral Testing
Through the Week Ending May 19, 2007 (Week 20)*

111100	Positive Influenza Total Number Specimens Submitted				
Week 20	0	2			
Season-to-Date	74	142			

^{*}Number positives also represent past weeks specimens submitted.

Antigenic Characterization of Missouri Influenza Isolates submitted to CDC by the State Public Health Laboratory: CDC antigenically characterizes a sample of positive Missouri influenza isolates, submitted through the Missouri Department of Health and Senior Services (DHSS), State Public Health Laboratory (SPHL). DHSS has submitted seventeen influenza isolates this season to CDC for antigenic characterization.

Results Received from CDC: CDC has antigenically characterized four of those isolates so far this season: two as influenza A/Hawaii/15/2001-like (H1N1) viruses and two as influenza A/New Caledonia/20/99-like (H1N1) viruses.

Isolates were characterized by hemagglutination-inhibition using post-infection ferret antisera and are related antigenically to A/Hawaii/15/2001, a minor antigenic variant of A/New Caledonia/20/99. A/New Caledonia/20/99 is the WHO recommended H1 component of the 2006-2007 vaccine for the Northern Hemisphere and 2007 vaccine for the Southern Hemisphere.

FDA's Vaccines and Related Biological Products Advisory Committee (VRBPAC) recommended that influenza vaccines to be used in the 2006-2007 season in the U.S. contain the following:

- An A/New Caledonia/20/99 (H1N1)-like virus;
- An A/Wisconsin/67/2005 (H3N2)-like virus (A/Wisconsin/67/2005 and A/Hiroshima/52/2005strains); and
- A B/Malaysia/2506/2004-like virus (B/Malaysia/2506/2004 and B/Ohio/1/2005 strains)

The influenza vaccine composition used in the 2006-2007 season in the U.S. is identical to that recommended by the World Health Organization on February 15, 2006.

Clusters/Outbreaks of Influenza-like Illness: No school closings were reported for week 20; no school closings were reported for last season. Thirteen school closings due to Influenza-like illness have been reported season-to-date, as compared to four school closings reported during this same time last season. No outbreaks have been reported this season as compared to nine last season.

Data Sources: Laboratory-confirmed cases are reported to DHSS through the passive communicable disease surveillance system. Suspected influenza clusters and outbreaks are reported through the active surveillance system. Pneumonia and influenza deaths are reported through the DHSS Bureau of Vital Records. Influenzalike illness data by age category and total number of patient visits by week are reported voluntarily by participants in the U.S. Influenza Sentinel Physicians Surveillance Network.

Find Us on the Web

This report may also be found on the DHSS Internet at: www.dhss.state.mo.us/Influenza/index.html.

National influenza surveillance information is available from the Centers for Disease Control and Prevention at: www.cdc.gov/ncidod/diseases/flu/weekly.htm.

Contact Us

The Missouri Department of Health and Senior Services after hour's number for reporting disease cases and emergencies is **1-800-392-0272**.

¹All data in this report are provisional and may change as reports are updated.

²Influenza activity codes are reported to CDC each Monday.